

## Supporting information

### MRI Visible Drug Eluting Magnetic Microspheres for Transcatheter Intra-Arterial Delivery to Liver

#### Tumors


Dong-Hyun Kim<sup>a\*</sup>, Jeane Chen<sup>b</sup>, Reed A. Omary<sup>c</sup>, Andrew C. Larson<sup>d</sup>

<sup>a</sup>Department of Radiology, Northwestern University, Chicago, IL 60611, USA; Robert H. Lurie Comprehensive Cancer Center, Chicago, IL 60611, USA.

<sup>b</sup>Department of Radiology, Northwestern University, Chicago, IL 60611, USA.

<sup>c</sup>Department of Radiology and Radiological Sciences, Vanderbilt University Medical Center, Nashville, TN, USA.

<sup>d</sup>Department of Bioengineering, University of Illinois at Chicago, Chicago, IL, USA; Department of Electrical Engineering and Computer Science, Evanston, IL, USA; International Institute of Nanotechnology (IIN), Northwestern University, Evanston, IL, USA; Robert H. Lurie Comprehensive Cancer Center, Chicago, IL 60611, USA. Department of Radiology, Northwestern University, Chicago, IL 60611, USA.



**Fig. S1.** Size variation in mean diameter of aqueous phase microspheres as a function of flow rate,  $Q_o=3\sim 40$  ul/min and  $Q_a=2$  and  $5$  ul/min.