

# Online Supporting Information for

## Drug-Bearing Supramolecular Filament Hydrogels as Anti-Inflammatory Agents.

Zhipeng Chen,<sup>a,b,d+</sup> Lei Xing,<sup>c+</sup> Qin Fan,<sup>a,b+</sup> Andrew G. Cheetham,<sup>d</sup> Ran Lin,<sup>d</sup> Barbara Holt,<sup>d</sup> Liwen Chen,<sup>d</sup> Yanyu Xiao,<sup>c\*</sup> and Honggang Cui.<sup>d,e,f\*</sup>

[+] These authors contributed equally to this work

<sup>a</sup> *Jiangsu Collaborative Innovation Center of Chinese Medicinal Resources Industrialization, Nanjing University of Chinese Medicine, Nanjing 210023, China*

<sup>b</sup> *Department of Pharmacy, Nanjing University of Chinese Medicine, Nanjing 210023, China*


<sup>c</sup> *State Key Laboratory of Natural Medicines, China Pharmaceutical University, Nanjing 210009, China*

<sup>d</sup> *Department of Chemical and Biomolecular Engineering and Institute for NanoBioTechnology, The Johns Hopkins University, Baltimore, MD 21218, USA*


<sup>e</sup> *Department of Oncology and Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA*

<sup>f</sup> *Center for Nanomedicine, The Wilmer Eye Institute, Johns Hopkins University School of Medicine, 400 North Broadway, Baltimore, MD 21231, USA*

[\*] Corresponding authors: [cpuyanyuxiao@163.com](mailto:cpuyanyuxiao@163.com); [hcu6@jhu.edu](mailto:hcu6@jhu.edu)



**Figure S1.** Representative mass spectrum of a Ket-VEVE conjugate (L-VEVE-(S)-Ket). All conjugates possess the same molecular mass.



**Figure S2.** *In vitro* dose–response relationship study of (S)-Ket and the (S)-Ket-conjugates against rabbit chondrocytes. The cells were incubated with the appropriate (S)-Ket and (S)-Ket-conjugate for 72 h and cell viability was determined by MTT assay. Data are given as mean  $\pm$  s.d. (n = 3).