

Supplementary materials

Dual-targeting Theranostic System with Mimicking Apoptosis to Promote Myocardial Infarction Repair *via* Modulation of Macrophages

Jingli Chen^{1,2,*}, Jun Yang^{1,*}, Ruiyuan Liu¹, Chenmeng Qiao¹, Zhiguo Lu¹, Yuanjie Shi¹, Zhanming Fan^{3,✉}, Zhenzhong Zhang^{2, ✉}, Xin Zhang^{1,✉}

1. State Key Laboratory of Biochemical Engineering, Institute of Process Engineering, Chinese Academy of Sciences, Beijing 100190, PR China

2. School of Pharmaceutical Sciences, Zhengzhou University, Zhengzhou 450001, PR China

3. Department of Radiology, Beijing An Zhen Hospital, Capital Medical University, Beijing Institute of Heart, Lung and Blood Vessel Disease, Beijing 100029, PR China

Corresponding author: xzhang@ipe.ac.cn, zhangzz08@126.com or fanzm120@126.com




Figure S1. Iron biodistribution *in vivo*. The concentration of iron in organs with various MIONs formulations after 1 day (d) and 1 month (m). Data are shown mean \pm SD. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.005$.




Figure S2. Histological and morphometric analysis. The Prussian blue staining of heart, liver, spleen, lung and kidney of different MIONs formulations 1 month after *i.v.* administration. 200× magnification. M: magnet.