

MicroRNA Biogenesis is Enhanced by Liposome-encapsulated Pin1 Inhibitor in Hepatocellular Carcinoma

Dan Sun,^{1,#} Shuangyan Tan,^{1,#} Yanli Xiong,^{1,2} Wenchen Pu,¹ Jiao Li,¹ Wei Wei,² Canhua Huang,^{1,3} Yu-Quan Wei,¹ Yong Peng^{1,*}

¹ *State Key Laboratory of Biotherapy and Cancer Center, National Clinical Research Center for Geriatrics, West China Hospital, Sichuan University, Chengdu, 610041, Sichuan, China.*

² *Key Laboratory of Bio-Resource and Eco-Environment of Ministry of Education, College of Life Sciences, Sichuan University, Chengdu 610065, China.*

³ *West China School of Basic Medical Sciences & Forensic Medicine, Sichuan University, Chengdu 610041, China.*

Supplementary Information




Figure S1. MTT assays after API-1 (blue) and API-LP (red) treatment of SK-Hep1 cells. Data are presented as mean \pm SD, n=3.




Figure S2. Immunoblotting analyses of XPO5, p-XPO5, ERK, p-ERK, and GAPDH levels in SK-Hep1 cells treated with DMSO, API-1 or API-LP (1 μ M).