

Erratum

Redox Regulation of Stem-like Cells Through the CD44v-xCT Axis in Colorectal Cancer: Mechanisms and Therapeutic Implications: Erratum

Huai-Qiang Ju^{1,2*}, Yun-Xin Lu^{1*}, Dong-Liang Chen^{1*}, Tian Tian¹, Hai-Yu Mo¹, Xiao-Li Wei¹, Jian-Wei Liao¹, Feng Wang¹, Zhao-Lei Zeng¹, Helene Pelicano³, Mitzi Aguilar², Wei-Hua Jia¹, Rui-Hua Xu¹✉

1. Sun Yat-sen University Cancer Center, State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, Guangzhou, 510060, China.
2. Department of Molecular and Cellular Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.
3. Department of Translational Molecular Pathology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.

*These three authors contributed equally to this work.

✉ Corresponding author: Prof. Rui-Hua Xu, M.D. Ph.D.; Address: 651 Dongfeng East Road, Guangzhou 510060, China; Tel: +86-20-8734-3228; Fax: +86-20-8734-3392; E-mail: xurh@sysucc.org.cn.

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>). See <http://ivyspring.com/terms> for full terms and conditions.

Published: 2025.01.09

Corrected article: *Theranostics* 2016; 6(8): 1160-1175. doi: 10.7150/thno.14848.

The authors regret some incorrect representative images were accidentally displayed during data organization, including flow cytometry scatter diagram in Figure 1F, Figure 4B, Figure S5B and colony image in Figure 6B. The authors confirm that these corrections do not change the result interpretation or conclusions of the article. The authors apologize for any inconvenience that the errors may have caused.

Corrected Figure 1F




Figure 1. (F) Reversion of H₂O₂-induced decreases in SP cells following NAC treatment. LoVo or SW620 cells were treated with 3 mM NAC for 2 h, followed by 100 µM H₂O₂ for 24 h.

Corrected Figure 4B




Figure 4. (B) CRC SP cell quantification in cells transfected with negative control (NC) or miR-1297 mimic.

Corrected Figure 6B




Figure 6. (B) LoVo and SW620 cells were incubated with 5FU, PEITC, or both for 2 weeks, and the cell colonies were fixed in formalin, stained with crystal violet, and counted.

Corrected Figure S5B




Figure A. Correct image for Figure S5B. (B) CRC cells were pretreated with SSZ (100 µM) for 48 h and then washed and cultured in fresh medium without the drug for 48 h to allow cell death to occur. Viable cells were harvested and then stained with Hoechst 33342 to identify SP cells, and the viable SP cells were quantified.