

## Erratum

# MAP4K3/GLK inhibits Treg differentiation by direct phosphorylating IKK $\beta$ and inducing IKK $\beta$ -mediated FoxO1 nuclear export and Foxp3 downregulation: Erratum

Jyun-Ni Chi<sup>1,§</sup>, Jhih-Yu Yang<sup>1,§</sup>, Chia-Hsin Hsueh<sup>1,§</sup>, Ching-Yi Tsai<sup>1</sup>, Huai-Chia Chuang<sup>1,✉</sup>, and Tse-Hua Tan<sup>1,2,✉</sup>

1. Immunology Research Center, National Health Research Institutes, Zhunan, Taiwan.

2. Department of Pathology & Immunology, Baylor College of Medicine, Houston, Texas, USA.

§These authors contributed equally to this work.

✉ Corresponding authors: Huai-Chia Chuang, 35 Keyan Road, Zhunan 35053, Taiwan. Phone: +886-37-206-166 ext. 37612; E-mail address: cinth@nhri.edu.tw. Tse-Hua Tan, 35 Keyan Road, Zhunan 35053, Taiwan. Phone: +886-37-206-166 ext. 37601; E-mail address: ttan@nhri.edu.tw


© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>). See <https://ivyspring.com/terms> for full terms and conditions.

Published: 2026.01.30


Corrected article: *Theranostics* 2022; 12(13): 5744-5760. doi: 10.7150/thno.72148.

The authors regret that the original version of our paper, unfortunately, contained an unnecessary IKK $\beta$  blot in Figure 3C right panel. The purified IKK $\beta$  protein was purchased from SignalChem and was used for all Figure 3C experiments. The blot of IKK $\beta$  input control should be shown only once to avoid misunderstanding. The reorganized version of Figure 3C is shown below. In addition, individual blots of Figure 3A left panel were inadvertently flipped upside down. The correct version of Figure 3A is shown below.

The corrections made in this erratum do not affect the original data and conclusions. The authors apologize for any inconvenience that the errors may have caused.



**Figure A. Corrected Figure 3C** Purified Flag-tagged FoxO1 and GST-tagged IKK $\beta$  proteins were used for *in vitro* binding assay. The relative protein levels of the protein complexes determined by densitometry analysis were shown in the bottom of panels.



**Figure B. Corrected Figure 3A** Co-immunoprecipitation experiments of Flag-tagged FoxO1 and HA-tagged IKK $\beta$  using lysates of HEK293T cells. IB, immunoblotting.